

Quantum Phase Estimation

Estimate eigenvalues of unitary operators
(without even diagonalizing the unitary)

Provable exponential/polynomial speedups

- Factoring - Shor's Algorithm
- solving linear systems of eq. - HHL
- Eigenvalues of molecular Hamiltonians

Estimating eigenvalues

why is it a phase?

Let M be a unitary / Hermitian matrix (Hermitian)
($M = M^\dagger$)

$$\begin{cases} MM^\dagger = I = M^\dagger M & (\text{unitary}) \\ MM^\dagger = M^2 = M^\dagger M & (\text{Hermitian}) \end{cases}$$

They are normal matrices $MM^\dagger = M^\dagger M$.

Then spectral decomposition states that
 $\exists$ orthonormal basis $|\psi_k\rangle$ s.t.:

$$M = \sum_k \lambda_k |\psi_k\rangle \langle \psi_k|$$

$$M = \sum_K \lambda_K (|\phi_K\rangle \langle \psi_K|)$$


$$\begin{aligned} M |\psi_j\rangle &= \sum_K \lambda_K (|\phi_K\rangle \langle \psi_K|) |\psi_j\rangle \\ &= \lambda_j |\psi_j\rangle \end{aligned}$$

$$M |\psi_j\rangle = \lambda_j |\psi_j\rangle$$

Moreover, unitary operators must preserve norm of vectors:

$$\|U|\psi_j\rangle\| = \|\lambda_j |\psi_j\rangle\| = |\lambda_j| \| |\psi_j\rangle \|$$

must $|\lambda| = 1$
lie on the unit circle



$$\lambda = \cos(\phi) + i \sin \phi$$

$$= e^{i\phi}$$

ϕ is 2π periodic $\lambda = e^{2\pi i \theta}$

$$U |\psi_j\rangle = e^{2\pi i \theta} |\psi_j\rangle$$

QPE estimates θ in binary floating point representation.

Let $U_{2^n \times 2^n}$ be unitary and K be the # bits
for the approximation:

QPE:

1) initialize $|0\rangle^n |0\rangle^K$

2) encode eigenstate $|\psi\rangle |0\rangle^K$

3) Apply operation U s.t:

$$| \psi_j \rangle | 0 \rangle^k \mapsto | \psi_j \rangle \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i \theta_j} | j \rangle$$

3) Apply operation U s.t:

$$| \psi_j \rangle | 0 \rangle^{\otimes n} \mapsto | \psi_j \rangle \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i \theta_j j} | j \rangle$$

Can we prepare it having only access to U ?

$$|0\rangle \xrightarrow{\text{QFT}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i \theta \cdot j} |j\rangle$$

$$|0\rangle \xrightarrow{\text{QFT}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i \theta j} |j\rangle$$



 QFT^{-1}

Apply inverse gives us the k -bit approximation $|\bar{\theta}\rangle$

4) Apply inverse QFT

Quantum phase estimation:

Given unitary $U_{2^n \times 2^n}$, eigenstate $|\psi\rangle$, k bits.

→ prepare $|0\rangle^n |0\rangle^k$

→ encode eigenstate $|\psi\rangle |0\rangle^k$

→ prepare operator O s.t

$$O |\psi\rangle |0\rangle^k \mapsto |\psi\rangle \frac{1}{\sqrt{2^n}} \sum_j e^{2\pi i \theta_j} |j\rangle$$

→ Apply inverse QFT to k bit register

$$\text{QFT}^{-1} \frac{1}{\sqrt{2^n}} \sum_j e^{2\pi i \theta_j} |j\rangle \mapsto |\tilde{\theta}\rangle$$

Encode the operation


0

k-bits for approximation

$$|0\rangle \xrightarrow{K} \boxed{H^{\otimes K}} \longrightarrow \frac{1}{\sqrt{2^K}} \sum_{\theta} |0\rangle$$


$$|\varphi\rangle \longrightarrow$$

$$U' |\varphi\rangle |0\rangle \longrightarrow e^{2\pi i \theta \cdot 0} |\varphi\rangle |0\rangle$$



Problem: $O(2^K)$ multicontrolled operations.

let's imagine that $|5\rangle = |101\rangle = |5\rangle$



Generalization: For qubit q in range $\{0, \dots, K-1\}$
 Apply $C-U^{2^q}$

Quantum Fourier Transform
has a product state
decomposition!

check Nielsen and Chuang Page 218

$$|j\rangle \rightarrow \frac{1}{2^{n/2}} \sum_{k=0}^{2^n-1} e^{2\pi i j k / 2^n} |k\rangle \quad (5.5)$$

$$= \frac{1}{2^{n/2}} \sum_{k_1=0}^1 \dots \sum_{k_n=0}^1 e^{2\pi i j (\sum_{l=1}^n k_l 2^{-l})} |k_1 \dots k_n\rangle \quad (5.6)$$

$$= \frac{1}{2^{n/2}} \sum_{k_1=0}^1 \dots \sum_{k_n=0}^1 \bigotimes_{l=1}^n e^{2\pi i j k_l 2^{-l}} |k_l\rangle \quad (5.7)$$

$$= \frac{1}{2^{n/2}} \bigotimes_{l=1}^n \left[\sum_{k_l=0}^1 e^{2\pi i j k_l 2^{-l}} |k_l\rangle \right] \quad (5.8)$$

$$= \frac{1}{2^{n/2}} \bigotimes_{l=1}^n \left[|0\rangle + e^{2\pi i j 2^{-l}} |1\rangle \right] \quad (5.9)$$

$$= \frac{\left(|0\rangle + e^{2\pi i 0 \cdot j_n} |1\rangle \right) \left(|0\rangle + e^{2\pi i 0 \cdot j_{n-1} j_n} |1\rangle \right) \dots \left(|0\rangle + e^{2\pi i 0 \cdot j_1 j_2 \dots j_n} |1\rangle \right)}{2^{n/2}}. \quad (5.10)$$

The product representation (5.4) makes it easy to derive an efficient circuit for the quantum Fourier transform. Such a circuit is shown in Figure 5.1. The gate R_k denotes the unitary transformation

$$R_k \equiv \begin{bmatrix} 1 & 0 \\ 0 & e^{2\pi i / 2^k} \end{bmatrix}. \quad (5.11)$$

$$|j\rangle \rightarrow \frac{1}{2^{n/2}} \sum_{k=0}^{2^n-1} e^{2\pi i j k / 2^n} |k\rangle \quad (5.5)$$

$$= \frac{1}{2^{n/2}} \sum_{k_1=0}^1 \dots \sum_{k_n=0}^1 e^{2\pi i j (\sum_{l=1}^n k_l 2^{-l})} |k_1 \dots k_n\rangle \quad (5.6)$$

$$= \frac{1}{2^{n/2}} \sum_{k_1=0}^1 \dots \sum_{k_n=0}^1 \bigotimes_{l=1}^n e^{2\pi i j k_l 2^{-l}} |k_l\rangle \quad (5.7)$$

$$= \frac{1}{2^{n/2}} \bigotimes_{l=1}^n \left[\sum_{k_l=0}^1 e^{2\pi i j k_l 2^{-l}} |k_l\rangle \right] \quad (5.8)$$


$$= \frac{1}{2^{n/2}} \bigotimes_{l=1}^n \left[|0\rangle + e^{2\pi i j 2^{-l}} |1\rangle \right] \quad (5.9)$$

$$= \frac{\left(|0\rangle + e^{2\pi i 0 \cdot j_n} |1\rangle \right) \left(|0\rangle + e^{2\pi i 0 \cdot j_{n-1} j_n} |1\rangle \right) \dots \left(|0\rangle + e^{2\pi i 0 \cdot j_1 j_2 \dots j_n} |1\rangle \right)}{2^{n/2}} \quad (5.10)$$

The product representation (5.4) makes it easy to derive an efficient circuit for the quantum Fourier transform. Such a circuit is shown in Figure 5.1. The gate R_k denotes the unitary transformation

$$R_k \equiv \begin{bmatrix} 1 & 0 \\ 0 & e^{2\pi i / 2^k} \end{bmatrix}. \quad (5.11)$$

H + C-R_k
For every control bit







Figure 5.1. Efficient circuit for the quantum Fourier transform. This circuit is easily derived from the product representation (5.4) for the quantum Fourier transform. Not shown are swap gates at the end of the circuit which reverse the order of the qubits, or normalization factors of $1/\sqrt{2}$ in the output.

Use pennylane's

qml.QFT (wires = estimation_wires)

Phase estimation full circuit.



⊗ for a given K ,

$$P(\tilde{\theta}) \geq 4/\pi^2 > 40\%$$

⊗ accuracy ϵ and Prob $> 1-\delta$

$$K > \log\left(\frac{1}{\epsilon} \left(\frac{1}{2\delta} + 2\right)\right)$$

CAVEATS

How to implement both

- U and $C U^2$
- eigenstate $|\psi\rangle$

Exercise : Consider

$$U = \begin{pmatrix} 1 & 0 \\ 0 & e^{2\pi i/5} \end{pmatrix}$$

angle
we want
to estimate
 $\phi = 1/5 = 0.2$

- qml. phaseshift (angle, wire)
- eigenstate?